

The Pause Between Breaths

Learning to WAIT in Recovery

Experience, Strength & Hope · The Recovery Files

A powerful exploration of reactive communication in recovery—how automatic responses persist long after substances are removed.

RELATIONSHIPS & COMMUNICATION · COMMUNICATION · INTERMEDIATE

The sound of shattering glass pulled me from sleep—a sound I knew intimately from my drinking days. Half-awake, I stumbled into the kitchen to find my partner, Elena, standing over broken fragments of what had been her grandmother's serving dish. Before my eyes had fully focused, before my brain had fully engaged, the words were already leaving my mouth:

"Jesus Christ, it's not even 7 AM. How the hell did you manage that?"

I watched her face collapse, watched how my words landed like another blow on top of her obvious distress. In that moment, twenty-six months into sobriety, I realized with absolute clarity that while I hadn't touched alcohol in over two years, I was still operating with the reactive brain of addiction.

My counselor introduced me to the concept of WAIT—What Am I Thinking? Why Am I Talking?—during my third month in outpatient treatment. At the time, I nodded politely while internally dismissing it as just another recovery acronym, less relevant than addressing the physical dependence that had dominated my life. I was there to stop drinking, not to analyze my communication style.

But that morning in the kitchen, standing amid fragments of glass and the fragments of connection broken by my thoughtless reaction, the concept suddenly seemed less like an optional recovery tool and more like an essential life skill I had never fully developed.

In active addiction, my responses were governed almost entirely by immediate emotional states—primarily irritation, defensiveness, and the constant low-grade anxiety of maintaining functional alcoholism. Substances provided a chemical buffer

between stimulus and response, temporarily dampening the intensity of emotions that might otherwise require more sophisticated management. When something triggered me—a comment from a colleague, a request from my partner, an unexpected change of plans—I had two primary strategies: react immediately with whatever emotion arose, or drink to suppress the discomfort of the trigger.

This reactive pattern became so ingrained that it survived well into sobriety. I had removed alcohol from my system but was still operating with neural pathways shaped by years of impulsive responses and chemical emotion management. The kitchen incident wasn't an isolated event but merely the most obvious example of a pattern that affected nearly all my interactions—with family, coworkers, friends, and even strangers. I was sober but not recovered in any meaningful relational sense.

The first component of WAIT—What Am I Thinking?—initially struck me as obvious to the point of uselessness. Of course I knew what I was thinking; they were my thoughts, weren't they? But when my therapist challenged me to actually articulate my thoughts in the moments before reactive responses, I discovered something unsettling: most of the time, I had no conscious awareness of the actual thought content driving my reactions.

Through dedicated practice, I began identifying the automatic thoughts that consistently preceded reactive outbursts. When Elena dropped the dish, the thoughts racing beneath my consciousness included: "This is going to make us late." "Now I'll have to clean this up." "She's so careless." "This day is already ruined." None of these thoughts were particularly accurate or helpful, but they created an emotional response that launched words from my mouth before I'd consciously processed the situation.

Learning to catch these thoughts required developing what my counselor called an "observing mind"—a part of consciousness that could notice thoughts arising without immediately identifying with or acting on them. This wasn't a natural skill for me; it required consistent practice through daily meditation and regular check-ins throughout the day, particularly during emotionally charged situations.

The second half of the WAIT acronym—Why Am I Talking?—proved even more challenging. When asked to identify my purpose in speaking during confrontational moments, my honest initial answer was usually some variation of "to release emotional pressure" or "to make the other person understand they're wrong." These motivations, while truthful, rarely aligned with my deeper values or the kind of person I wanted to be in recovery.

With practice, I began asking myself better questions before speaking: "Will these words strengthen or damage this relationship?" "Am I trying to connect or control?" "Is this communication in service of my recovery values or my ego?" "What need am I trying to meet by speaking right now, and is there a more effective way to meet that need?"

My sponsor framed it this way: "Getting sober creates the possibility of recovery. Practices like WAIT are what actually build recovery, one intentional response at a time." This distinction—between the absence of substances and the presence of recovery—helped me understand why many people, myself included, could maintain physical sobriety while still experiencing significant dysfunction in relationships and emotional management.

Disengaging from arguments proved particularly challenging early in this practice. My partner and I had established patterns where tension would escalate rapidly from minor irritations to major conflicts, with neither of us having effective off-ramps from the emotional intensity. Learning to say "I need some time to process this before responding" felt awkward and artificial at first, both to express and to hear.

Developing specific language for temporary disengagement helped immensely: "I want to have this conversation in a way that's productive, but I'm feeling too reactive right now. Can we pause for thirty minutes so I can gather my thoughts?" This approach acknowledged the importance of the issue while creating space for more intentional engagement.

Physical practices became essential supports for these mental and emotional shifts. Deep breathing exercises, particularly box breathing (inhaling for four counts, holding for four, exhaling for four, holding for four), provided immediate physiological intervention during triggering situations. This technique directly countered the shallow, rapid breathing pattern that accompanied emotional reactivity, sending signals to my nervous system that the perceived emergency wasn't actually life-threatening.

The most profound shift occurred when I began recognizing that most of my reactive responses were fundamentally about fear rather than anger. Behind the quick irritation when Elena dropped the dish were fears about control, competence, judgment, and ultimately worthiness of love and belonging. These deeper fears had driven both my

addictive behaviors and my communication patterns, creating a self-reinforcing cycle where substances temporarily numbed the fear while reactive communication consistently reinforced it.

A particularly meaningful milestone occurred approximately eighteen months into consistent WAIT practice. During a heated discussion about financial priorities, I noticed the familiar physical sensations of reactivity arising—increased heart rate, shallow breathing, muscle tension in my jaw and shoulders. Instead of being hijacked by these sensations, I observed them with genuine curiosity, asked myself what thoughts were generating them, and chose a response aligned with my deeper values rather than immediate emotional release.

Today, thirty-five months sober and about two years into intentional WAIT practice, the changes are substantial though still evolving. Reactive responses haven't disappeared entirely but have diminished significantly in both frequency and intensity. The pause between stimulus and response has expanded from microseconds to meaningful moments of choice. Communication, particularly during emotionally charged situations, more consistently reflects my actual values rather than conditioned reactivity.

This morning, Elena dropped a mug while making coffee. The sound triggered a momentary surge of irritation—the physiological response still occurs—but the space between that sensation and my response has grown large enough to accommodate conscious choice. What am I thinking? "Accidents happen; this isn't a big deal." Why am I talking? "To connect and offer support." My response flowed from this awareness: "You okay? Don't worry about the mess—I'll grab the broom while you finish making coffee."

The WAIT practice hasn't eliminated conflict from our relationship or my other interactions. Disagreements, misunderstandings, and differences of opinion remain part of human connection. But these inevitable frictions now occur within a container of greater awareness and intentionality, allowing them to become opportunities for growth and deeper understanding rather than triggers for reactive damage.

I've come to understand that recovery isn't just about abstaining from substances but about developing the capacity to be fully present—with ourselves, with others, with the full range of human experience—without needing to escape or react impulsively. The WAIT practice supports this broader recovery by strengthening the muscle of conscious choice in moments previously governed by automatic reactivity. The pause between stimulus and

response, between breath and breath—this is where recovery lives, one WAIT at a time.

THE MATCHED SELF-ASSESSMENT

The Pause Between Breaths: Reactive Communication in Recovery

This self-assessment is designed to help individuals in recovery evaluate their patterns of reactive communication and develop skills for more intentional responses. The assessment explores both reactive patterns that often persist beyond substance use and the development of mindful communication practices that support sustainable recovery.

Instructions

For each statement, rate how often it applies to you using the scale:

- 1 - Never: This doesn't apply to you
- 2 - Rarely: This occasionally applies
- 3 - Sometimes: This sometimes applies
- 4 - Often: This frequently applies
- 5 - Always: This consistently applies

Part 1: Reactive Communication Patterns

1. I respond to situations with immediate emotional reactions before thinking
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
2. I say things in the heat of the moment that I later regret
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
3. I find myself speaking to release emotional pressure rather than to communicate effectively
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
4. My communication style changes noticeably when I'm stressed, tired, or triggered
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
5. I have difficulty pausing before responding in emotional situations
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
6. I speak without clear awareness of what I'm thinking or why I'm talking
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
7. My first response to mistakes or accidents is irritation or criticism
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
8. I escalate quickly from minor disagreements to major conflicts
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
9. I feel physically activated (racing heart, muscle tension, shallow breathing) during challenging conversations
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
10. I communicate in ways that damage connection rather than strengthen it when upset
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always

Part 2: Mindful Communication Skills

11. I can identify my automatic thoughts before responding to triggering situations
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
12. I practice pausing between stimulus and response in challenging moments
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
13. I ask myself "What am I thinking?" when I notice emotional reactivity arising
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
14. I consider "Why am I talking?" before speaking in important conversations
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
15. I can disengage temporarily from heated interactions to gather my thoughts
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
16. I notice physical signs of reactivity (breathing changes, tension) before they control my responses
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
17. I communicate in ways that align with my deeper values rather than immediate emotions
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
18. I consider how my words will impact relationships before speaking
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
19. I can differentiate between primary emotions (fear, hurt) and secondary reactions (anger, irritation)
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always
20. I take responsibility promptly when my communication has been reactive or harmful
☐ 1 Never ☐ 2 Rarely ☐ 3 Sometimes ☐ 4 Often ☐ 5 Always

Reflection Questions

Understanding Reactive Patterns

1. In what specific situations or with which people do you find yourself most reactive?

2. What physical sensations do you experience when becoming reactive (tension, breathing changes, etc.)?

3. What automatic thoughts typically precede your reactive communications?

4. How did substances previously help you manage reactive tendencies?

Communication in Recovery

5. How have your communication patterns changed since achieving sobriety?

6. What impact does your communication style have on your recovery journey?

7. How does your current communication affect your relationships with family, friends, and colleagues?

8. What emotions typically drive your reactive responses (fear, shame, insecurity, etc.)?

The WAIT Practice

9. What challenges do you face in implementing "What Am I Thinking?" before responding?

10. What are your common answers to "Why Am I Talking?" in difficult conversations?

11. How could creating more pauses in your communication benefit your recovery?

12. What specific language could help you disengage temporarily from heated interactions?

Physical Dimensions

13. What breathing practices help you interrupt the physiological aspects of reactivity?

14. How does physical self-care (sleep, nutrition, exercise) affect your communication patterns?

15. What bodily cues could serve as early warning signs of impending reactivity?

16. What grounding techniques help you return to presence during triggering conversations?

Growth Opportunities

17. What specific communication skills would you most like to develop further?

18. What accountability systems could support your practice of more mindful communication?

19. How might improved communication reduce potential relapse triggers?

20. What daily practices could help you strengthen the pause between stimulus and response?

Your scores, and what they mean, are in the app.