

High Risk Situations in Recovery

Relapse Prevention

Step-Working Guide · The Recovery Files

In this workbook

1. **High Risk Situations in Recovery** — key ideas, reading, worksheet
Each topic's scored self-assessment and mind map are in the app.

TOPIC

High Risk Situations in Recovery

KEY IDEAS

Understanding High Risk Situations

High risk situations are those dangerous moments when substance use begins to appear reasonable, logical, or inevitable despite previously established recovery commitments.

The Deceptive Nature

These situations represent not simply external triggers but complex internal processes where addictive thinking actively prepares the groundwork for relapse while simultaneously dismantling protective barriers.

What begins as seemingly logical reasoning gradually evolves into a self-reinforcing cycle of isolation, exposure to triggers, and disconnection from recovery supports.

The 12 Dimensions of High Risk

High risk situations manifest across twelve interconnected dimensions:

Relief-Seeking Rationalization

Substances as solution

Comparative Reasoning

Using seems better

Control Illusion

Believing in moderation

Secrecy Justification

Hidden use doesn't count

Recovery Restart Fantasy

Can get sober again

Support Network Disconnection

Withdrawing from help

High-Risk Association

Seeking using contacts

Environmental Testing

Exposing to triggers

Help Rejection

No one can help

Surrender to Inevitability

Relapse is certain

Last Threshold Awareness

Imminent risk

Recovery Abandonment

Discarding recovery

Relief-Seeking Rationalization

Relief-seeking rationalization involves the belief that substances would provide relief from emotional pain, solve current problems, or improve one's situation.

Common Manifestations

- Thinking substances would provide relief from emotional pain
- Believing using would solve current problems
- Seeing substances as medicine for psychological suffering
- Romanticizing immediate effects while minimizing consequences
- Experiencing emotional states as unbearable without chemical relief

Recovery Responses

- Utilize healthy coping skills to address emotional pain
- Recognize emotional pain as temporary and manageable
- Seek appropriate help for managing intense emotional states
- Maintain perspective that substances ultimately worsen pain

Control Illusion & Secrecy Justification

Control Illusion

Convincing oneself that controlled use is possible:

- Convincing yourself you could use without losing control
- Believing recovery knowledge enables moderate use
- Creating detailed scenarios about controlled use
- Selectively remembering controlled use, forgetting loss of control

Secrecy Justification

Believing hidden use "doesn't count":

- Believing undisclosed use "doesn't really count"
- Planning to hide evidence of substance use
- Thinking secret use would have fewer consequences
- Believing recovery appearances could be maintained

Network Disconnection & High-Risk Association

Support Network Disconnection

Withdrawing from recovery-supportive relationships:

- Withdrawing from recovery-supportive relationships
- Hiding true thoughts and feelings from supports
- Becoming unreachable to people who know recovery well
- Creating excuses to avoid recovery meetings

High-Risk Association

Seeking influences that facilitate relapse:

- Spending time with people who actively use
- Seeking relationships with enablers
- Maintaining connections with former using network
- Distancing from recovery-committed people

Environmental Testing & Help Rejection

Environmental Testing

Creating opportunities for relapse through exposure:

- Deliberately placing yourself where substances are present
- Testing ability to resist by exposure to triggers
- Visiting locations associated with past use
- Removing environmental safeguards

Help Rejection

Eliminating sources of intervention:

- Believing no one could help with current struggle
- Deciding not to reach out before potential relapse
- Rejecting the idea that support could help
- Believing mind is already made up about using

Surrender, Last Threshold & Abandonment

Surrender to Inevitability

- Feeling relapse is inevitable regardless of actions
- Experiencing powerlessness to prevent using
- Viewing situation through lens of predetermined failure

Last Threshold Awareness

- Recognizing imminent risk of relapse
- Experiencing dangerous calm about impending use
- Making specific plans for how and when to use

Recovery Abandonment

- Believing recovery principles no longer apply
- Rejecting previously accepted recovery aspects
- Discarding identification as person in recovery

The 12 Recovery Responses

For each vulnerability, there's a corresponding recovery response:

Relief-Seeking→Pain Management Alternatives

Comparative Reasoning→Reality-Based Comparison

Control Illusion→Control Reality Acceptance

Secrecy Justification→Honesty Commitment

Restart Fantasy→Recovery Restart Reality

Network Disconnection→Support Network Activation

High-Risk Association→Protective Association

Environmental Testing→Environmental Protection

Help Rejection→Help Receptivity

Surrender to Inevitability→Choice Empowerment

Last Threshold→Crisis Response Protocol

Recovery Abandonment→Recovery Recommitment

Building Relapse Prevention Resilience

The path forward isn't about eliminating challenging thoughts—it's about developing awareness to recognize high risk thinking and tools to implement effective responses.

Key Prevention Practices

- Regular Risk Assessment: Weekly examination of thoughts and behaviors
- Staged Response Protocol: Tiered responses for different risk levels
- Personalized Trigger Management: Strategies for your unique triggers
- Emergency Response Planning: Concrete plans for imminent risk
- Accountability Structure: Multiple layers of support and oversight

"Between the emergence of high risk thinking and your response lies a space of choice that grows larger with each day of recovery practice."

CORE READING

Dancing with Danger: How I Orchestrated My Own Relapse

Creating the Conditions for Return to Use

The descent toward relapse began when my emotional pain reached an unbearable threshold. "It can't get any worse than feeling this way," became my internal mantra, a thought that appeared rational but actually represented addiction's masterful distortion of reality.

I convinced myself that the emotional suffering I was experiencing constituted a unique emergency that justified extraordinary measures. This dangerous rationalization—that relapse was inevitable and perhaps even necessary—represented the first significant step toward return to use.

The allure of chemical relief grew more compelling as I fixated on memories of how substances had once seemingly solved my problems. I began romanticizing my relationship with alcohol and drugs, selectively remembering the temporary escape while conveniently forgetting the devastating consequences that inevitably followed.

The Illusion of Control

The most sophisticated deception my addiction created was the belief that I could use again in a controlled manner. "I've learned so much in recovery," I reasoned. Despite constructing this elaborate justification, a quiet voice of truth persisted—I knew with absolute certainty that controlled use was impossible for me.

To further weaken my recovery commitment, I created artificial distinctions between public and private behavior, convincing myself that a relapse only "counted" if others knew about it. This compartmentalization represented a profound misunderstanding of how recovery works.

As these distorted thoughts intensified, I systematically disconnected from my recovery network, eliminating the people who might have challenged these dangerous rationalizations. This deliberate isolation represented addiction's classic strategy of isolating the prey from the protection of the herd.

As relapse grew closer, I began engaging in increasingly risky behaviors that practically guaranteed eventual use. I would drive past old using locations "just to see what's changed." I would sit in bars ordering non-alcoholic drinks, testing my resilience while surrounded by triggers.

No Lifelines Remaining

By the time actual substance use occurred, I had created a situation where I had no lifelines remaining and, more disturbingly, no desire to reach for them even if they existed. My mind was made up, the decision already finalized in some fundamental way that made external input seem irrelevant.

Returning to treatment has meant confronting these patterns with painful clarity. I now understand that high-risk situations aren't just external circumstances I encounter but conditions I actively create through progressively distorted thinking and deliberate choices. By sharing these patterns with others in treatment, I not only maintain accountability but potentially help others recognize similar progressions before they culminate in relapse.

WORKSHEET

High Risk Situations Worksheet

Personal Mapping & Relapse Prevention Planning

MAPPING

Creating Your Personalized High Risk Map

For each area, reflect on where you've been and where you're heading.

Relief-Seeking→Alternatives-Healthy

Where I was: 'I thought substances would provide relief from emotional pain rather than utilizing healthy coping skills'

Where I am now:

Where I'm heading:

Control-Illusory→Acceptance-Reality

Where I was: 'I convinced myself I could use without losing control rather than acknowledging proven inability to control'

Where I am now:

Where I'm heading:

Secrecy-Justified→Honesty-Committed

Where I was: 'I believed hidden use didn't count rather than maintaining rigorous honesty'

Where I am now:

Where I'm heading:

Network-Disconnected→Support-Activated

Where I was: 'I withdrew from recovery relationships rather than deliberately increasing contact during vulnerability'

Where I am now:

Where I'm heading:

Environment-Testing→Protection-Environmental

Where I was: 'I deliberately placed myself where substances were present rather than avoiding exposure'

Where I am now:

Where I'm heading:

Inevitability-Surrendered→Choice-Empowered

Where I was: 'I felt relapse was inevitable rather than recognizing relapse as choice'

Where I am now:

Where I'm heading:

PATTERNS

Your Personal High Risk Patterns

List specific examples for each category that affect you personally.

Relief-Seeking Rationalization

Example: I believe alcohol would help me sleep during periods of insomnia

Comparative Reasoning

Example: I tell myself using 'just once' would be better than continuing to feel depressed

Control Illusion

Example: I fantasize about being able to drink only on special occasions

Secrecy Justification

Example: I think about how I could hide evidence of drinking from my partner

Recovery Restart Fantasy

Example: I tell myself I could easily get sober again after 'just one weekend' of using

Support Network Disconnection

Example: I stop returning my sponsor's calls when thinking about using

High-Risk Association

Example: I reconnect with former drinking buddies 'just to catch up'

Environmental Testing

Example: I deliberately walk through the liquor section at the grocery store

Help Rejection

Example: I decide in advance I won't call anyone if urges become overwhelming

Surrender to Inevitability

Example: I start to believe relapse is just a matter of time for me

Last Threshold Awareness

Example: I find myself checking store hours of nearby liquor stores

Recovery Abandonment

Example: I start to believe AA/NA principles don't apply to my unique situation

ACTION

Action Planning for Risk Management

Select 2-3 specific risk patterns to address.

Risk Pattern 1

Pattern to Address:

30-Day Goal:

Action Steps This Week:

Accountability Measures:

Risk Pattern 2

Pattern to Address:

30-Day Goal:

Action Steps This Week:

Accountability Measures:

Risk Pattern 3

Pattern to Address:

30-Day Goal:

Action Steps This Week:

Accountability Measures:

EMERGENCY

Emergency Response Protocol

Develop your concrete plan for when relapse feels imminent.

Early Warning Signs

What are your personal early warning signs that risk is escalating?

Immediate Steps

What are 3-5 specific actions you will take when you notice warning signs?

Emergency Contacts

Who are 3-5 people you will call, in order, during a crisis? Include names and numbers.

Safe Locations

Where can you go that is safe and substance-free during a crisis?

Last Threshold Protocol

What specific steps will you take if you're at the final moment before using?

Remember: Managing high risk situations isn't about eliminating all challenging thoughts—it's about developing sophisticated awareness and effective response strategies that recognize high risk patterns as deceptive internal processes requiring comprehensive intervention.

www.TheRecoveryFiles.com

Embrace, Empower & Evolve